

Integral Representations of (p, q) -Forms With Weighted Factors On Complex Manifolds*

Wu Xiaoqin

Li Lunhuan

(Department of Math., Jimei Teachers College, Xiamen) (Department of Math., Xiamen Univ.)

Abstract In this paper we construct a Cauchy-Leray kernel with weighted factors and obtain an integral representation of (p, q) -forms and a Koppelman-Leray-Norguet formula on a complex manifold.

Key words Weighted factor, integral representation, complex manifold.

0 Introduction

In the research of complex functions of several variables, integral representations hold a very important position. In 1970s, Henkin obtained the integral representations of the solution of $\bar{\partial}$ -equation. After that, integral representations are extensively being studied and applied.

In 1982, Andersson & Bendtsson[2] introduced the weighed factors in integral representations in C^n . By choosing appropriate factors, applications of integral representations are more extensive and convinient. In 1990, Andersson [3] obtained the integral representations of (p, q) -forms in a stein manifold.

In 1989, by using Chern-class theory, Berndtsson [1] obtained the integral representations of (p, q) -forms in a complex manifold. It is the most general integral representations so far.

This paper, as an extension of [1], the author constructs a Cauchy-Leray kernel with weighted factors and obtains an integral representation of (p, q) -forms with weighted factors and Koppelman-Leray-Norguet formula on a complex manifold.

1 Construction of the kernel

Let X be a complex manifold (for example, $M \times M$, M is a complex manifold of complex dimension n). Let Y be a complex submanifold of codimension p (for example, Δ , Δ is diag-

* Supported by Natureal Science Foundation of Fujian.

nal of $M \times M$). Suppose that $E \xrightarrow{\pi} X$ is a holomorphic vector—bundle of rank p , which has a smooth section η to E and $Y = \{\eta = 0\}$. Moreover

$$d\eta_1 \wedge d\eta_2 \cdots \wedge d\eta_p \wedge d\bar{\eta}_1 \wedge d\bar{\eta}_2 \wedge \cdots \wedge d\bar{\eta}_p \neq 0 \text{ on } Y$$

where η_i are the coefficients of η about some frame.

Let E^* the dual bundle of E and ξ be an any section of E^* . Let the connections of E and E^* be D and D^* , respectively. Let a local frame of E be e_1, \cdots, e_p , its duals be e_1^*, \cdots, e_p^* . Then

$$e_1^* \wedge e_1 \wedge \cdots \wedge e_p^* \wedge e_p$$

is independent of the choice of frame.

The exterior product on $E^* \oplus E$ can be extended to forms with values in this bundle. In the terms of our local trivialization a form with values in $\Lambda^{r,s}$ can be written

$$\alpha = \sum_{|I|=r, |J|=s} a_{IJ} e_I^* \wedge e_J$$

where a_{IJ} are forms, $I = (i_1, \cdots, i_r)$, $J = (j_1, \cdots, j_s)$. If

$$\beta = \sum b_{KL} e_K^* \wedge e_L$$

is another such object, we define:

$$\alpha \wedge \beta = \sum a_{IJ} \wedge b_{KL} e_I^* \wedge e_J \wedge e_K^* \wedge e_L$$

In other words, forms can commute with vectors from $E^* \oplus E$ and product between forms is the usual one.

Now, we define the differential d that is independent of the frame. Let

$$d\alpha = \sum_{|I|=r, |J|=s} da_{IJ} e_I^* \wedge e_J$$

Let $\beta = \sum b_{KL} e_K^* \wedge e_L$, and a_{IJ} be m -forms, then

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^m \alpha \wedge d\beta$$

Let $\theta = D^2$ be the curvature of E . Let

$$\det(I + \frac{it}{2\pi} \theta) = \sum_{k=0}^p C_k[\theta] t^k$$

In terms of Chern—Weil thorem we know that $C_k[\theta]$ is independent of D . Let $\theta = (\theta_{ij})_{p \times p}$ and

$\bar{\theta} = \sum_{i,j=1}^p \theta_{ij} e_i^* \wedge e_j$ We have formula

$$(\frac{i}{2\pi})^p \frac{1}{p!} \bar{\theta}^p = C_p[\theta] e_1^* \wedge e_1 \wedge \cdots e_p^* \wedge e_p$$

It is independent of frame. Let

$$\Lambda = e_1^* \wedge e_1 \wedge \cdots e_p^* \wedge e_p$$

Now, we construct Cauchy—Leray kernel with weighted factors. Let Q_0, Q_1, \cdots, Q_N are smooth sections of E^* . Then $\langle Q_j, \eta \rangle$ is a smooth function in $X \times X$, $j=0, 1, \cdots, N$, and

$$d \langle Q_j, \eta \rangle = \langle D^* Q_j, \eta \rangle + \langle Q_j, D\eta \rangle$$

is 1-form in $X \times X$, Let

$$M\Lambda = C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^{k-1} \sum_{|\alpha|=p-k} \frac{1}{\alpha!} g^{(\alpha)} (D^* Q_0 \wedge D\eta)^{\alpha_0} \wedge \cdots \wedge (D^* Q_N \wedge D\eta)^{\alpha_N} \wedge \bar{\theta}^k$$

where $\alpha_0 + \dots + \alpha_N = p - k$, $\alpha! = \alpha_0! \dots \alpha_N!$, $C_p = (2\pi i)^{-p}$, $g(\lambda_0, \dots, \lambda_N)$ is a holomorphic function, and

$$g^{(*)} = \left[\frac{\partial^0}{\partial \lambda_0^0} \dots \frac{\partial^N}{\partial \lambda_N^N} \right] g(< Q_0, \eta >, \dots, < Q_N, \eta >)$$

Let

$$B = -C_p \sum_{|\alpha|=p} \frac{1}{\alpha!} g^{(*)} (D^* Q_0 \wedge D\eta)^{\alpha_0} \wedge \dots \wedge (D^* Q_N \wedge D\eta)^{\alpha_N}$$

Similarly to [3], we have

$$\begin{aligned} dB &= -C_p \sum_{|\alpha|=p} \frac{1}{\alpha!} g^{(*)} d[(D^* Q_0 \wedge D\eta)^{\alpha_0} \wedge \dots \wedge (D^* Q_N \wedge D\eta)^{\alpha_N}] \\ &= -C_p \sum_{|\alpha|=p} \frac{1}{\alpha!} g^{(*)} \sum_{j=0}^N \alpha_j d(D^* Q_j \wedge D\eta) \wedge (D^* Q_0 \wedge D\eta)^{\alpha_0} \wedge \dots \\ &\quad \wedge (D^* Q_N \wedge D\eta)^{\alpha_{j-1}} \wedge \dots \wedge (D^* Q_N \wedge D\eta)^{\alpha_N} \end{aligned}$$

Let $N = dM$ Then we have $N\Lambda = (dM)\Lambda = d(M\Lambda)$

Now we replace g in $M\Lambda$ and $N\Lambda$ by $H(\lambda_0)g(\lambda_1, \dots, \lambda_N)$ and Q_0 by $t\xi$. we denote by M_t and N_t the forms so obtained when differentials are also taken with respect to t . we also use the short hand notation

$$(D^* Q \wedge D\eta)^* = (D^* Q_1 \wedge D\eta)^{\alpha_1} \wedge \dots \wedge (D^* Q_N \wedge D\eta)^{\alpha_N}$$

Thus

$$\begin{aligned} M_t \Lambda &= C_p \sum_{k=0}^p \binom{p}{k} (-1)^{k-1} \sum_{j+|\alpha|=p-k} \frac{1}{j! \alpha!} H^{(j)} g^{(*)} (dt\xi \wedge D\eta \\ &\quad + tD^* \xi \wedge D\eta)^j \wedge (D^* Q \wedge D\eta)^* \wedge \partial^k \\ N_t \Lambda &= -C_p \sum_{j+|\alpha|=p} \frac{1}{j! \alpha!} H^{(j)} g^{(*)} d[(dt\xi \wedge D\eta + tD^* \xi \wedge D\eta)^j \wedge (D^* Q \wedge D\eta)^*] \\ &\quad + C_p \sum_{k=1}^{p-1} \binom{p}{k} (-1)^{k-1} \sum_{j+|\alpha|=p-k} \frac{1}{j! \alpha!} \{ dH^{(j)} \wedge g^{(*)} (dt\xi \wedge D\eta + tD^* \xi \wedge D\eta)^j \\ &\quad \wedge (D^* Q \wedge D\eta)^* \wedge \partial^k \\ &\quad + H^{(j)} dg^{(*)} \wedge (dt\xi \wedge D\eta + tD^* \xi \wedge D\eta)^j \wedge (D^* Q \wedge D\eta)^* \wedge \partial^k \\ &\quad + H^{(j)} g^{(*)} d(dt\xi \wedge D\eta + tD^* \xi \wedge D\eta)^j \wedge (D^* Q \wedge D\eta)^* \wedge \partial^k \\ &\quad + H^{(j)} g^{(*)} (dt\xi \wedge D\eta + tD^* \xi \wedge D\eta)^j \wedge d[(D^* Q \wedge D\eta)^* \wedge \partial^k] \} \end{aligned}$$

Proposition 1.1 If M_t and N_t are as above, then

$$(d_t + d)M_t = N_t$$

where d is the differential which is independent of t .

Let $H(t) = \exp(-t)$. If $a \neq 0$ we define

$$\int_0^\infty \exp(-at) t^j dt = j! / a^{j+1}$$

This "integral" obeys such rules as differentialing under the integral sign and formula

$$\int_0^\infty \frac{d}{dt} x(t) dt = -x(0)$$

Definition 1.2 $K(\xi, \eta, Q, g) = \int_0^\infty M_t$ is called Cauchy-Lerey Kernal with weighted fac-

tors, where the integral on t is d_t in the most right side.

Let $R = \int_0^\infty N_t$ and $P = M_t|_{t=0}$. Thus we have

$$dK = \int_0^\infty dM_t = - \int_0^\infty d_t M_t + \int_0^\infty N_t = M_t|_{t=0} + R = P + R$$

where we let $\langle \xi, \eta \rangle = 0$, Now we concretely calculate K, P, R .

The term containing dt in $M_t \Lambda$ is

$$C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^k \sum_{j+|\alpha|+p-k} \frac{1}{j! \alpha!} H^{(j)} g^{(\alpha)} \xi \wedge D\eta (D^* \xi \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^* \wedge j t^{j-1} dt$$

Therefore,

$$K \Lambda = \int_0^\infty \mu_t \Lambda = C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^k \sum_{j+|\alpha|+p-k} \frac{1}{j! \alpha!} g^{(\alpha)} \frac{\xi \wedge D\eta (D^* \xi \wedge D\eta)^{j-1}}{\langle \xi, \eta \rangle^j} \wedge (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^* \quad (1.3)$$

Because $\xi \wedge \xi = 0$, so when we replace ξ in (1.3) by $\varphi \xi$ (1.3) is invariant.

$$P \Lambda = M_t|_{t=0} \Lambda = M_t \Lambda|_{t=0} \quad (1.4)$$

$$= C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^k \sum_{|\alpha|+p-k} \frac{1}{\alpha!} g^{(\alpha)} (D^* \xi \wedge D\eta)^* \wedge \bar{\theta}^* \text{ Let } \xi = \sum \xi_j e_j^*, \eta = \sum \eta_j e_j, D^*$$

$$\xi = \sum (D^* \xi)_j e_j^*, D\eta = \sum (D\eta)_j e_j,$$

$$C^* \xi = d(D^* \xi) = \sum d(D^* \xi)_j e_j^* \text{ and } c\eta = d(D\eta) = \sum d(D\eta)_j e_j.$$

Then the first term of $\{ \}$ in Nt which contains dt is

$$- H^{(j+1)} g^{(\alpha)} [\langle \xi, \eta \rangle (D^* \xi \wedge D\eta)^j - j (\langle D^* \xi, \eta \rangle + \langle \xi, D\eta \rangle) \wedge \xi \wedge D\eta \wedge (D^* \xi \wedge D\eta)^{j-1}] \wedge (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^* \wedge t^j dt$$

where

$$\begin{aligned} dH^{(j)} &= H^{(j+1)} d(-t \langle \xi, \eta \rangle) \\ &= -H^{(j+1)} [dt \langle \xi, \eta \rangle + t (\langle D^* \xi, \eta \rangle + \langle \xi, D\eta \rangle)] \end{aligned}$$

The second term of $\{ \}$ in $Nt \wedge$ which contains dt is

$$\begin{aligned} &- H^{(j)} \sum_{i=1}^N \frac{\partial}{\partial \lambda_i} g^{(\alpha)} (\langle D^* Q_i, \eta \rangle + \langle Q_i, D\eta \rangle) \wedge \xi \wedge D\eta \\ &\wedge (D^* \xi \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^* \wedge t^{j-1} dt \end{aligned}$$

The third term of $\{ \}$ in Nt which contains dt is

$$\begin{aligned} &H^{(j)} g^{(\alpha)} [(D^* \xi \wedge D\eta)^j - (d\xi \wedge D\eta + \xi \wedge c\eta) \wedge (D^* \xi \wedge D\eta)^{j-1} \\ &+ \xi \wedge D\eta \wedge (j-1) (C^* \xi \wedge D\eta - D^* \xi \wedge c\eta) \wedge (D^* \xi \wedge D\eta)^{j-2}] \\ &\wedge (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^* \wedge j t^{j-1} dt \end{aligned}$$

The fourth term of $\{ \}$ in Nt which contains dt is

$$\begin{aligned} &H^{(j)} g^{(\alpha)} \xi \wedge D\eta \wedge (D^* \xi \wedge D\eta)^{j-1} \wedge \left[\sum_{i=1}^N \alpha_i (C^* Q_i \wedge D\eta - D^* Q_i \wedge C\eta) \right. \\ &\left. \wedge (DQ \wedge D\eta)^{s_1 \cdots s_{j-1} \cdots s_N} \wedge \bar{\theta}^* + (D^* Q \wedge D\eta)^* \wedge \bar{\theta}^{j-1} \right] \end{aligned}$$

where

$$(DQ \wedge D\eta)^{s_1 \cdots s_{j-1} \cdots s_N} = (D^* Q_{i_1} \wedge D\eta)^{s_1 \cdots s_{j-1}} \wedge (D^* Q_{i_j} \wedge D\eta)^{s_j \cdots s_{j-1}} \cdots \wedge (D^* Q_{i_N} \wedge D\eta)^{s_N}$$

Therefore we have

$$\begin{aligned}
 R\Lambda &= \int_0^\infty N_t \\
 &= C_p \sum_{j+|\alpha|=\rho} \frac{1}{\alpha!} g^{(\alpha)} \left[\frac{d\xi \wedge D\eta \wedge (D\xi^* \wedge D\eta)^{j-1} \wedge (D^*Q \wedge D\eta)^*}{\langle \xi, \eta \rangle^j} \right. \\
 &\quad + \frac{\xi \wedge C\eta \wedge (D\xi^* \wedge D\eta)^{j-1} \wedge (D^*Q \wedge D\eta)^*}{\langle \xi, \eta \rangle^j} \\
 &\quad - \frac{\xi \wedge D\eta \wedge (j-1)(C^*\xi \wedge D\eta - D^*\xi \wedge c\eta) \wedge (D^*\xi \wedge D\eta)^{j-2} \wedge (D^*Q \wedge D\eta)^*}{\langle \xi, \eta \rangle^j} \\
 &\quad - \frac{\xi \wedge D\eta \wedge (D\xi^* \wedge D\eta)^{j-1}}{\langle \xi, \eta \rangle^j} \\
 &\quad \cdot \sum_{i=1}^N \alpha_i (C^*Q_i \wedge D\eta - D^*Q_i \wedge C\eta) (D^*Q \wedge D\eta)^{a_1 \cdots a_{j-1} \cdots a_i} \\
 &\quad - j \frac{(\langle D^*\xi, \eta \rangle + \langle \xi, D\eta \rangle) \wedge \xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{j-1} \wedge (D^*Q \wedge D\eta)^*}{\langle \xi, \eta \rangle^{j+1}} \Big] \\
 &\quad + C_p \sum_{k=1}^{p-1} \binom{p}{k} (-1)^{k-1} \sum_{j+|\alpha|=\rho-k} \frac{1}{j! \alpha!} \\
 &\quad \cdot \{ g^{(\alpha)}
 \end{aligned}$$

$$\begin{aligned}
 &\quad j(\langle D^*\xi, D\eta \rangle + \langle \xi, D\eta \rangle) \wedge \xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{j-1} - \langle \xi, \eta \rangle (D^*\xi \wedge D\eta)^{j-1} \\
 &\quad \wedge (D^*Q \wedge D\eta)^* \wedge \partial^k \\
 &\quad - \sum_{i=1}^N \frac{\partial}{\partial \lambda_i} g^{(\alpha)} (\langle D_i^*, \wedge D\eta \rangle + \langle Q_i, D\eta \rangle) \wedge \frac{\xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{j-1}}{\langle \xi, \eta \rangle^j} \wedge (D^*Q \wedge D\eta)^* \wedge \partial^k \\
 &\quad + g^{(\alpha)} \left[\frac{D^*\xi \wedge D\eta)^j - (d\xi \wedge D\eta + \xi \wedge (\eta) \wedge (D^*\xi \wedge D\eta)^{j-1}}{\langle \xi, \eta \rangle^j} \right. \\
 &\quad \left. + \frac{(j-1)\xi \wedge D\eta \wedge (C^*\xi \wedge D\eta - D\xi \wedge C\eta) \wedge (D^*\xi \wedge D\eta)^{j-2}}{\langle \xi, \eta \rangle^j} \right] \wedge (D^*Q \wedge D\eta)^* \\
 &\quad \wedge \partial^k \\
 &\quad + g^{(\alpha)} \frac{\xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{j-1}}{\langle \xi, \eta \rangle^j} \wedge \left[\sum_{i=1}^N \alpha_i (C^*Q_i \wedge D\eta - D^*Q_i \wedge C\eta) \right. \\
 &\quad \wedge (D^*Q \wedge D\eta)^{a_1 \cdots a_{j-1} \cdots a_N} \wedge \partial^k \\
 &\quad \left. + (D^*Q \wedge D\eta)^* \wedge kd\partial \wedge \partial^{k-1} \right] \}
 \end{aligned}$$

Espacially, if $g \equiv 1, Q_i \equiv 0, i=1, \cdots, N$, we get

$$K\Lambda = C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^k \frac{\xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{p-k-1}}{\langle \xi, \eta \rangle^{p-k}} \wedge \partial^k \quad (1.6)$$

This is the Cauchy-Leray kernel constructed by B. Berndtsson in [1]. And

$$P\Lambda \equiv 0, \quad \text{Thus} \quad P \equiv 0 \quad (1.7)$$

$$\begin{aligned}
 R\Lambda &= C_p \sum_{k=0}^{p-1} \binom{p}{k} (-1)^k \left[\frac{(d\xi \wedge D\eta \wedge \xi \wedge C\eta) \wedge (D^*\xi \wedge D\eta)^{p-k-1}}{\langle \xi, \eta \rangle^{p-k}} \wedge \partial^k \right. \\
 &\quad - \frac{(p-k-1)\xi \wedge D\eta \wedge (C^*\xi \wedge D\eta - D^*\xi \wedge C\eta) \wedge (D^*\xi \wedge D\eta)^{p-k-2}}{\langle \xi, \eta \rangle^{p-k}} \wedge \partial^k \\
 &\quad - \frac{(p-k)(\langle D^*\xi, \eta \rangle + \langle \xi, D\eta \rangle \wedge \xi \wedge D\eta) \wedge (D^*\xi \wedge D\eta)^{p-k-1}}{\langle \xi, \eta \rangle^{p-k-1}} \wedge \partial^k \\
 &\quad \left. + \frac{\xi \wedge D\eta \wedge (D^*\xi \wedge D\eta)^{p-k-1}}{\langle \xi, \eta \rangle^{p-k}} \wedge kd\partial \wedge \partial^{k-1} \right]
 \end{aligned} \quad (1.8)$$

2 Integral representation of (p,q) -forms with weighted factors

Now let M be a complex manifold and its complex dimension be n . Let $X=M \times M$ and E be a holomorphic vector bundle on X whose rank is n . Let η be a section of E so that

$$\Delta = \{\eta = 0\} = \{(\xi, z) \in M \times M, \xi = z\}$$

Definition 2.1 Let ξ be a smooth section of E^* . ξ is called allowable about η if for any $A \subset \subset X$ we have

$$|\xi| \leq C_A |\eta|, \quad |\langle \xi, \eta \rangle| \geq C_A |\eta|^2$$

where C_A and C_A are constants only related with A .

Let $g \equiv 1$. If ξ is allowable near Δ and K, P , and R are (1.6) (1.7) (1.8), respectively, then

$$dK = [\Delta] + R \quad (R = -C_A[\theta]) \quad (2.2)$$

where $[\Delta]$ denotes the current that integrations is over $\Delta \subset M \times M$. This is the result of [1]. If K, P and R are general, i. e. as in (1.3) (1.4) and (1.5), noting that the principal part of K is just as in (1.6) we get from (2.2) following theorem:

Theorem 2.3 If ξ is allowable about and $g(0)=1$, following formula is satisfied

$$dK = [\Delta] + R + P$$

Theorem 2.4 If $D \subset \subset M$ is a domain with $C^{(1)}$ -boundary and if f is any smooth k -form, we have the formula

$$f(z) = \int_D K \wedge df + d \int_D K \wedge f + (-1)^k \int_{\partial D} K \wedge f + \int_D P \wedge f + \int_D R \wedge f, z \in D$$

If E is a holomorphic vector bundle and η is its section, then for any $C^{(1)}(p,q)$ -form $f(z)$ in D we have

$$\begin{aligned} f(z) = & \int_D K_{p,q} \wedge \bar{\partial} f + \bar{\partial} \int_D K_{p,q-1} \wedge f + (-1)^{p+q} \int_{\partial D} K_{p,q} \wedge f \\ & + \int_D P_{p,q} \wedge f + \int_D R_{p,q} \wedge f \end{aligned}$$

where lower index (p,q) means the component which has bidegree (p,q) in z and $K_{p,-1} \equiv 0$

The proof of the theorem is standard and we omit it.

3 The Koppelman-Leray-Norguet formula with weighted factors

Definition 3.1 Let M be a complex manifold of complex dimension n . An open set $D \subset \subset M$ is called to have a piecewise $C^{(1)}$ -boundary if there exists a finite open covering $\{U_i\}_{i=1}^h$ of an open neighbourhood U of ∂D and $C^{(1)}$ -functions $\rho_j \rightarrow (1 \leq j \leq h)$ such that

- 1) $D \cap U = \{z \in U \mid \text{for } 1 \leq j \leq h, \text{ either } z \notin U, \text{ or } \rho_j(z) < 0\}$
- 2) for $1 \leq i_1 < \dots < i_l \leq h$, the 1-forms $d\rho_{i_1}, \dots, d\rho_{i_l}$

are linearly independent over R at every point of $\bigcap_{v=1}^1 U_{iv}$. And $\{U_j, \rho_i\}$ is called a frame of D .

For every ordered subset $I = \{i_1, \dots, i_l\}$ of $\{1, 2, \dots, h\}$ define

$$S_I = \{z \in \partial D \cap (\bigcap_{i \in I} U_i) : \rho_i(x) = 0, i \in I\}$$

and choose the orientation on S_I such that the orientation is skew symmetric in the component of I and follwing two equation hold when D is given the natural orientation:

$$\partial D = \sum_{j=1}^h S_j, \partial S_I = \sum_{j=1}^h S_{Ij}$$

Let

$$\sigma = \{\lambda = (\lambda_0, \dots, \lambda_h) \in R^{h+1}, \lambda_j \geq 0, \sum_{j=0}^h \lambda_j = 1\}$$

be the standard h -simplex in R^{h+1} with cononical orientation, for every ordered subset $J = \{j_1, \dots, j_m\}$ of $\{0, \dots, h\}$ with strictly increasing components, set

$$\sigma_J = \{\lambda \in \sigma : \sum_{j \in J} \lambda_j = 1\}$$

The orientation of each σ_j is chosen so that

$$\partial \sigma_J = \sum_{v=1}^m (-1)^{v-1} \sigma_{j_1 \dots j_{v-1} j_{v+1} \dots j_m}$$

where \hat{j}_v means that j_v is omitted. With this orientation we have

$$\partial(S_I \times \sigma_J) = \partial S_I \times \sigma_J + (-1)^{|I|} S_I \times \partial \sigma_J$$

Lemma 3. 2

$$\partial(\sum_I (-1)^{|I|} S_I \times \sigma_{0I}) = \sum_I S_I \times \sigma_I - \partial D \times \sigma_0$$

where the summation is over all ordered subsets I of $\{1, \dots, h\}$ with strictly increasing components, and $0I = \{0, i_1, \dots, i_l\}$

Let ξ, ξ_1, \dots, ξ_h be sections of E^* which are allowable about η . Let g and Q_i be as in ξ_i . Let

$$t^*(\xi, \xi_1, \dots, \xi_h, \eta, \lambda) = \sum_{i=1}^h \lambda_i \frac{\xi_i}{\langle \xi_i, \eta \rangle} + \lambda_0 \frac{\xi}{\langle \xi, \eta \rangle}$$

and

$$\begin{aligned} & \bar{K}(\xi, \xi_1, \dots, \xi_h, \eta, Q, g, \lambda) \Lambda \\ &= C_n \sum_{k=0}^{n-1} \binom{n}{k} (-1)^k \sum_{j+|a|=n-k} \frac{1}{\alpha!} g^{(a)} t^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^a \wedge \theta^k \end{aligned}$$

Lemma 3. 3 $\bar{K}(\xi, \xi_1, \dots, \xi_h, \eta, Q, g, \lambda) |_{\sigma_{0I}} = \bar{K}(\xi_1, \dots, \xi_h, \eta, Q, g, \lambda)$

$$\bar{K}(\xi, \xi_1, \dots, \xi_h, \eta, Q, g, \lambda) |_{\sigma_0} = \bar{K}(\xi, \eta, Q, g)$$

Lemma 3. 4 $d\bar{K}(\xi, \xi_1, \dots, \xi_h, \eta, Q, g, \lambda)$

$$\begin{aligned} &= (d_{\xi, \eta} + d_\lambda) \bar{K}(\xi, \xi_1, \dots, \xi_h, \eta, Q, g, \lambda) \\ &= C_n \sum_{j+|a|=n} \frac{1}{\alpha!} g^{(a)} [dt^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^a \\ &\quad + t^* \wedge C\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^a \\ &\quad - t^* \wedge D\eta \wedge (j-1)(C^* t^* \wedge D\eta - D^* t^* \wedge C\eta) \wedge (D^* t^* \wedge D\eta)^{j-2} \\ &\quad \wedge (D^* Q \wedge D\eta)^a \\ &\quad - t^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge \sum_{i=1}^N \alpha_i (C^* Q_i \wedge D\eta - D^* Q_i \wedge C\eta) \end{aligned}$$

$$\begin{aligned}
& \wedge (D^* Q \wedge D\eta)^{\sigma_1 \cdots \sigma_{l-1} \cdots \sigma_N}] \\
& + C_{\sigma} \sum_{k=1}^{n-1} \binom{n}{k} (-1)^k \sum_{j+|\sigma|=n-k} \frac{1}{\alpha!} \left\{ \sum_{l=1}^N \frac{\partial}{\partial \lambda_l} g^{(\alpha)}(< D^* Q_l, \eta > + < Q_l, D\eta >) \right. \\
& \quad \wedge t^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^{\sigma} \wedge \bar{\theta}^k \\
& \quad + g^{(\alpha)} dt^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^{\sigma} \wedge \bar{\theta}^k \\
& \quad + g^{(\alpha)} t^* \wedge C\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^{\sigma} \wedge \bar{\theta}^k \\
& \quad - g^{(\alpha)} t^* \wedge D\eta \wedge j-1 (C^* t^* \wedge D\eta - D^* t^* \wedge C\eta) \wedge (D^* t^* \wedge D\eta)^{j-2} \\
& \quad \wedge (D^* Q \wedge D\eta)^{\sigma} \wedge \bar{\theta}^k \\
& \quad - g^{(\alpha)} t^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge \sum_{l=1}^N \alpha_l (C^* Q_l \wedge D\eta - D^* Q_l \wedge C\eta) \\
& \quad \wedge (D^* Q \wedge D\eta)^{\sigma_1 \cdots \sigma_{l-1} \cdots \sigma_N} \wedge \bar{\theta}^k \\
& \quad - g^{(\alpha)} t^* \wedge D\eta \wedge (D^* t^* \wedge D\eta)^{j-1} \wedge (D^* Q \wedge D\eta)^{\sigma} \wedge kd\bar{\theta} \wedge \bar{\theta}^{k-1} \\
& \triangleq T(\xi, \xi_1, \cdots, \xi_h, \eta, Q, g, \lambda) \wedge
\end{aligned}$$

The proofs of Lemma 3.3 and Lemma 3.4 are simple and are omitted.

Theorem 3.5 Let $D \subset \subset M$ and its boundary ∂D be $C^{(1)}$ -piecewise. Let $\xi, \xi_1, \cdots, \xi_h$ be sections of E^* that are allowable about η . If f is a (p, q) -form whose coefficients are in $C^{(1)}(D)$, then we have following formula:

$$\begin{aligned}
f(z) &= (-1)^{p+q} \sum_I \int_{S_I \times \sigma_I} \bar{K}_{p,q} \wedge f + (-1)^{p+q+1} \sum_I (-1)^{(I)} \int_{S_I \times \sigma_{OI}} \bar{T}_{p,q} \wedge f \\
&\quad - \sum_I (-1)^{|I|} \int_{S_I \times \sigma_{OI}} \bar{K}_{p,q} \wedge \bar{\partial} f + \int_D \bar{K}_{p,q} \wedge \bar{\partial} f + \bar{\partial}_z \int_D \bar{K}_{p,q-1} \wedge f \\
&\quad + \int_D R_{p,q} \wedge f + \int_D P_{p,q} \wedge f, \quad z \in D
\end{aligned}$$

The proof of Theorem 3.5 Applying Theorem 2.4 and Stokes formula, we have

$$\begin{aligned}
\int_{\partial D \times \sigma_0} \bar{K} \wedge f &= \sum_I \int_{S_I \times \sigma_I} \bar{K} \wedge f - \sum_I (-1)^{|I|} \int_{S_I \times \sigma_{OI}} d(\bar{K} \wedge f) \\
&= \sum_I \int_{S_I \times \sigma_I} \bar{K} \wedge f - \sum_I (-1)^{|I|} \int_{S_I \times \sigma_{OI}} \bar{T} \wedge f \\
&\quad + (-1)^{p+q-1} \sum_I (-1)^{|I|} \int_{S_I \times \sigma_{OI}} \bar{K} \wedge \bar{\partial} f
\end{aligned}$$

Therefore the theorem is proved.

Reference

- 1 Berndtsson B. Cauchy-Leray forms and vector bundles, Department of Math, Chalmers Univ. of Technology, The University of Goteborg, Sweden. 1989.
- 2 Andersson M. Weighted Solution Formulas for the $\bar{\partial}$ -equation on domains with piecewise smooth strictly pseudocomplex boundaries, Math. Ann. 1973, 206
- 3 Andersson M. Weighted Solution Formulas for the $\bar{\partial}$ -equation on a Stein Manifold, preprint 1989.
- 4 Range R M & Siu Y T. Uniform estimates for the $\bar{\partial}$ -equation on domains with piecewise smooth strictly pseudoconvex boundaries, Math. Ann. 1973, 206
- 5 Zhong Tongde. Singular Integrals and Integral Representations in Several Complex Variables, Contemporary Math. 1993, 4421

(下转第70页)

Math. Anal. Appl. ,1992,165:346~

- 7 Ladas G. Recent developments in the oscillations of delay difference equations, in "Differential equations: stability and control", Dekker, New York, 1990.
- 8 Gedrgion D A etal. Oscillation of neutral difference equations, Appl. Anal. ,Nos. 3-4, 1989, 33:243~253
- 9 Lalli, B S etal, On oscillations and existence of positive solutions of neutral difference equations, J. Math. Anal. Appl. , 1991, 158:213~233
- 10 Lalli B S and Zhang B G. On existence of positive solutions and bounded oscillations for neutral difference equations, J. Math. Anal. Appl. , 1992, 166:272~287

On The Second Order Neutral Delay Difference Equations with Variable Coefficients

Shen Jianhua

(Hunan University, Changsha, 410082)

Abstract In this paper, we are concerned with the oscillation and nonoscillation of the second order neutral delay difference equation

$$\Delta^2(X_n - C_n X_{n-m}) = p_n X_{n-k}, n \geq n_0 \quad (*)$$

where, C_n, p_n are real numbers, m, k, n_0 are given nonnegative integers such that $p_n \geq 0, p_n \neq 0, n \geq n_0$ and $m \geq 1$. $\Delta X_n = X_{n+1} - X_n$. We show that if $C_n \geq 0$ then E_q . (*) has a unbounded positive solution. Sufficient and also necessary and sufficient conditions are obtained for oscillation of all bounded solutions of E_q . (*).

Key Word Neutral difference equation, oscillation, nonoscillation

(上接第 59 页)

复流形上 (p, q) 形式的带权因子的积分表示

吴晓勤

(厦门集美师范专科学校数学系, 厦门 36000)

李轮换

(厦门大学数学系, 厦门 361005)

摘 要 本文在[1]的基础上, 通过构造带权的 Cauchy-Leray 核, 得到了一般复流形上的 (p, q) 形式的带权因子的积分表示和带权因子的 Koppelman-Leray-Noryuet 公式.

关键词 权因子, 积分表示, 复流形